

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
26 August 2004 (26.08.2004)

PCT

(10) International Publication Number  
**WO 2004/071400 A3**

(51) International Patent Classification<sup>7</sup>: **C07D 213/56**,  
C07C 233/51, 237/40, A61K 31/4406, 31/165, A61P  
17/06, 35/00

(21) International Application Number:  
PCT/IB2004/000401

(22) International Filing Date: 9 February 2004 (09.02.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/447,915 14 February 2003 (14.02.2003) US  
10/770,035 2 February 2004 (02.02.2004) US

(71) Applicant: **SHENZHEN CHIPSCREEN BIO-SCIENCES LTD.** [CN/CN]; Research Institute of Tsinghua University, Suite C301, P.O. Box 28, High-Tech Industrial Park Nanshan District, Shenzhen, Guangdong 518057 (CN).



(72) Inventors: **LU, Xian-Ping**; Research Institute of Tsinghua University, Suite C301, P.O. Box 28, High-Tech Industrial Park Nanshan District, Shenzhen, Guangdong 518057

(CN). **LI, Zhibin**; Research Institute of Tsinghua University, Suite C301, P.O. Box 28, High-Tech Industrial Park Nanshan District, Shenzhen, Guangdong 518057 (CN). **XIE, Aihua**; Research Institute of Tsinghua University, Suite C301, P.O. Box 28, High-Tech Industrial Park Nanshan District, Shenzhen, Guangdong 518057 (CN). **LI, Boyu**; Research Institute of Tsinghua University, Suite C301, P.O. Box 28, High-Tech Industrial Park Nanshan District, Shenzhen, Guangdong 518057 (CN). **NING, Zhiqiang**; Research Institute of Tsinghua University, Suite C301, P.O. Box 28, High-Tech Industrial Park Nanshan District, Shenzhen, Guangdong 518057 (CN). **SHAN, Song**; Research Institute of Tsinghua University, Suite C301, P.O. Box 28, High-Tech Industrial Park Nanshan District, Shenzhen, Guangdong 518057 (CN). **DENG, Tuo**; Research Institute of Tsinghua University, Suite C301, P.O. Box 28, High-Tech Industrial Park Nanshan District, Shenzhen, Guangdong 518057 (CN). **HU, Weiming**; Research Institute of Tsinghua University, Suite C301, P.O. Box 28, High-Tech Industrial Park Nanshan District, Shenzhen, Guangdong 518057 (CN).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

[Continued on next page]

(54) Title: HISTONE DEACETYLASE INHIBITORS OF NOVEL BENZAMIDE DERIVATIVES WITH POTENT DIFFERENTIATION AND ANTI-PROLIFERATION ACTIVITY



(57) Abstract: The present invention is related to the preparation and pharmaceutical use of novel benzamide derivatives as defined in the specification of formula (I) as histone deacetylase inhibitors (HDACI), their preparations and the methods of using these compounds or their pharmaceutically acceptable salt in the treatment of cell proliferative diseases, e.g. cancer and psoriasis.

WO 2004/071400 A3



AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

**(84) Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK,

**(88) Date of publication of the international search report:**

16 June 2005

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

**HISTONE DEACETYLASE INHIBITORS OF NOVEL BENZAMIDE  
DERIVATIVES WITH POTENT DIFFERENTIATION AND  
ANTI-PROLIFERATION ACTIVITY**

5

Priority is indicated herein from U.S. Provisional Application Ser. No. 60/447,915, filed February 14, 2003.

**FIELD OF THE INVENTION**

10 The present invention relates to the preparation and use of novel benzamide derivatives as histone deacetylase inhibitors for treatments of differentiation- and/or proliferation-related disorders such as cancer and psoriasis.

**BACKGROUND OF THE INVENTION**

15 Aberrant expression of genes plays significant roles in the pathogenesis or pathological alterations in cancer, endocrine-related disorders, immune/inflammatory diseases, genetic diseases, and neurological diseases. The human genome is packaged into chromatin that consists of DNA, histones and non-histone proteins. Chromatin structure is an important factor in determining whether a particular gene is expressed or not. In  
20 general, condensed chromatin mediates transcriptional repression, whereas transcriptionally active genes are in areas of open chromatin. Nucleosomes form the basic repeating unit of chromatin, and consist of DNA wrapped around a histone octomer that is formed by four histone partners, namely an H3-H4 tetramer and two H2A-H2B dimmers. Histone H1 acts like a linker to stabilize the higher-order folding  
25 by electrostatic neutralization of the linker DNA segments through a positively charged carboxy-terminal domain. Therefore, the dynamic higher-order structure of nucleosomes defines distinct levels of chromatin organization and consequently, gene activation. Ricky W. Johnstone, "Histone deacetylase inhibitors: novel drugs for the treatment of cancer", *Nature Reviews Drug Discovery* 2002, 1: 287. The capacity of

histone octamer to compact DNA is influenced by a number of post-translational modifications that occur on the N-terminal histone tails. One modification involves the reversible acetylation and deacetylation of the epsilon-amino group of lysine moieties found within the histone tails. The net level of acetylation of N-terminal histone tails is controlled by the activities of two families of enzymes, the histone acetyltransferases (HATs) and histone deacetylases (HDACs). In addition to HATs and HDACs, other factors also participate in determination of chromatin structure, including methyl-CpG-binding protein and adenosine triphosphate-dependent chromatin-remodeling complexes that can directly recruit HDACs, which leads to repression of gene activation (see review *Current Opinion in Oncology* 2001, 13:477-483).

The identification of coactivator complexes that possess intrinsic HAT activity strongly supports the connection between histone acetylation and transcriptional activation. Similarly, transcriptional repressor complexes have been shown to recruit HDACs to the promoter of target genes (*Bioassays* 1998, 20:615). Sequence-specific transcription factors, such as nuclear hormone receptor superfamily, cyclic adenosine 3', 5'-monophosphate-related enhancer binding protein (CREB), and signal transducer and activator of transcription-1 (STAT-1), interact with distinct coactivators and corepressors inside the complex of transcriptional machinery in a DNA-context and tissue-context selective fashion, resulting in selective regulation of gene expression networks. These regulatory networks govern homeostasis of our body's physiological functions and perturbation of those networks causes disorders and/or profoundly affects progression of diseases. Therefore, modulation of complex interactions of transcription machinery provides novel intervention strategies for the treatments of cancer, endocrine-related disorders, immune/inflammatory diseases, genetic diseases, and neurodegeneration (Korzus, E., *et al.*, Transcription Factor-specific Requirements for

Coactivator and Their Acetyltransferase Functions. *Science* 1998, 279: 703-707; McKenna, N.J. and B.W. O'Malley, Combinatorial Control of Gene Expression by Nuclear Receptors and Coregulators. *Cell* 2002, 108(4):465-474; Pazin, M.J. and J.T. Kadonaga, What's Up and Down with Histone Deacetylation and Transcription? *Cell* 1997, 89(3):325-328; Zhong, H., R.E. Voll, and S. Ghosh, Phosphorylation of NF-B p65 by PKA Stimulates Transcriptional Activity by Promoting a Novel Bivalent Interaction with the Coactivator CBP/p300. *Molecular Cell* 1998, 1(5):661-671; Steffan JS. *et al.*, Histone deacetylase inhibitors arrest polyglutamine-dependent neurodegeneration in *Drosophila*, *Nature* 2001, 413:691-694; HDAC inhibitor VX-563 from Vertex Pharmaceuticals proceeds for genetic disorders, 2002 EDGAR online News, US20020115716A1, WO0056153A1).

For instance, cell development and differentiation are governed by the hierarchical order of sequential gene activation, which is controlled at the level of chromatin structure. Genetic alterations or mutations that cause constitutive activation of oncoproteins such as RAS, or inactivation of tumor suppressors such as p53, affect a myriad of molecular programs, including transcription. In addition, genetic abnormalities that result in improper targeting of HATs and HDACs to certain *loci*, functional inactivation of HATs, overexpression of HDACs or epigenetic changes due to DNA hyper- and hypomethylation, can shift balance between cell development and differential programs that often lead to tumor onset and progression (see review *Current Opinion Genet. Development* 1999, 9: 40-48 and 175-184). Several human cancers have been associated with malfunctions in HAT and HDAC activity. One example is the translocation of chromosomes 15 and 17 seen in the majority of acute promyelocytic leukemia (APL) patients. In APL, chromosomal translocations produce fusion proteins that contain RARalpha, PML (promyelocytic leukaemia protein), and PLZF (promyelocytic zinc finger). These aberrant proteins bind to retinoic acid

response elements, recruit HDACs with high affinity through enhanced binding for SMRT corepressor and are not responsive to retinoids, resulting in the constitutive repression of RAR-targeted genes (*Oncogene* 2001, 20:7204-7215). Retinoid acid receptor (RAR) is a ligand-activated transcriptional modulator that is important for myeloid differentiation. RAR heterodimerized with its partner RXR binds to retinoid acid response element, located in promoter region of target genes, and in the absence of retinoids, represses transcription by recruiting SIN3/HDAC through NCOR and SMRT corepressors. Addition of ligand releases the HDAC complexes from RAR/RXR, and allows subsequent binding of HATs, such as TIF2 and CBP, to activate transcription.

Therefore, the coordinated activation and repression of genes that contain functional retinoid acid response elements is essential to myeloid cell differentiation. Furthermore, addition of HDAC inhibitors can restore sensitivity of APL cells to retinoid-induced myeloid cell differentiation, indicating that aberrant histone deacetylation is a key process in leukaemogenesis.

There are reports that histone deacetylases, when overexpressed, silence the expression of tumor suppressor genes that are natural brakes against tumor growth. For example, p53, a critical regulator of cell proliferation, transmits signals to genes that control the cell cycle and apoptosis when cells are under stress. The functions are principally controlled by the ability of p53 to bind to DNA with sequence-specificity and to activate transcription. Inactivation of this property of p53, mostly by mutations that occur in the central DNA-binding domain, often leads to malignancy. It has been demonstrated that CBP/p300 can up-regulate p53 through core histone acetylation and p53 acetylation. (W. Gu and R.G. Roeder, Activation of p53 Sequence-Specific DNA Binding by Acetylation of the p53 C-Terminal Domain. *Cell* 1997, 90(4): 595-606.) Conversely, mammalian HDAC-1, HDAC-2, and HDAC-3 are shown to be capable of downregulating p53 function by deacetylation of both core histone and p53 (Juan, L.-J.,

et al., Histone Deacetylases Specifically Down-regulate p53-dependent Gene Activation. *The Journal of Biological Chemistry* 2000, 275(27): 20436-20443).

These data show that inappropriate transcriptional repression mediated by HDACs is a  
5 common molecular mechanism that is used by oncoproteins, and alterations in  
chromatin structure can impinge on normal cellular differentiation, which leads to  
tumor formation and other hyperproliferative disorders. Therefore, the inhibition of  
HDAC activity seems to be a rational therapeutic pathway for cancers and other  
hyperproliferative diseases.

10

Several classes of HDAC inhibitors have been identified, including (1) short-chain fatty  
acids, e.g. butyrate and phenylbutyrate; (2) organic hydroxamic acids, e.g.  
suberoylanilide hydroxamic acid (SAHA) and trichostatin A (TSA); (3) cyclic  
tetrapeptides containing a 2-amino-8-oxo 9,10-expoxydecanoyl (AOE) moiety, e.g.  
15 trapoxin and HC-toxin; (4) cyclic peptides without the AOE moiety, e.g. apicidin and  
FK228; and (5) benzamides, e.g. MS-275 (EP0847992A1, US2002/0103192A1,  
WO02/26696A1, WO01/70675A2, WO01/18171A2).

Butyric acid acts as an inhibitor of cell proliferation and an inducer of  
20 cytodifferentiation due primarily to its activity of inhibiting histone deacetylase. (A.  
Nudelman and A. Rephaeli, Novel Mutual Prodrug of Retinoic and Butyric Acids with  
Enhanced Anticancer Activity. *J. Med. Chem.* 2000, 43(15): 2962-2966.)  
Phenylbutyrate has been used as a single agent in the treatment of  $\beta$ -thalassemia,  
toxoplasmosis, and malaria. It is also reported to be successful in treating refractory  
25 APL in combination with RA (retinoid acid). (R.P. Warrell. et al., Therapeutic targeting  
of transcription in acute promyelocytic leukemia by use of an inhibitor of histone  
deacetylase. *J. Natl. Cancer Inst.* 1998, 90(21): 1621-1625.) Another fatty acid,

valproic acid, which is a potent anticonvulsant, mood stabilizer and teratogen, is also a direct inhibitor of histone deacetylase. (C.J. Phiel et al., Histone Deacetylase Is a Direct Target of Valproic Acid, a Potent Anticonvulsant, Mood Stabilizer, and Teratogen. *The Journal of Biological Chemistry* 2001, 276(39): 36734-36741; EP1170008A1).

5

A set of benzamides was discovered to have HDAC inhibitory activity in the low micromolar range. A lead compound, MS-275, from this set of benzamides is being tested by Mitsui Chemicals, Inc. and it is the first HDAC inhibitor to demonstrate oral anticancer activity in animal models with no severe side effects. (A. Saito et al., A synthetic inhibitor of histone deacetylase, MS-27-275, with marked *in vivo* antitumor activity against human tumors. *Proceedings of the National Academy of Sciences of the United States of America* 1999, 96(8): 4592-4597; EP 0847992 A1). MS-275 is currently under clinical trials in the University of Maryland Greenebaum Cancer Center for leukemia patients and by the U.S. National Cancer Institute for advanced solid tumors. (E.B. Levit, Clinical Trials in Leukemia focus on New Treatment Approaches. 2001 Release - University of Maryland Medical News 2001 Maryland <http://www.umm.edu/news/releases/karp.html>, A Phase I Study of an Oral Histone Deacetylase Inhibitor, MS-275, in Refractory Solid Tumors and Lymphomas. 2001, National Cancer Institute). However, there is still a need to discover new compounds with improved profiles, such as stronger HDAC inhibitory activity.

15

20

#### SUMMARY OF THE INVENTION

The current invention provides compounds that exhibit differentiation-inducing and proliferation-inhibiting effects and are useful as therapeutic treatment or improving agents for disorders related to differentiation and/or proliferation, such as cancer and psoriasis. In particular, they are highly effective against hematological malignancy and solid carcinoma.

25



## BRIEF DESCRIPTION OF THE FIGURES


Figure 1 graphically demonstrates the transcriptional activation of various nuclear hormone receptors by example HDAC inhibitors, i.e., Trichostatin A, MS-275 as well  
5 as an example compound of the present invention.

## DETAILED DESCRIPTION OF THE INVENTION

Various publications are cited throughout the present application. The contents of these publications and contents of documents cited in these publications are herein  
10 incorporated herein by reference.

The present invention provides compounds represented by formula (I), or its stereoisomer, enantiomer, diastereomer, hydrate, or pharmaceutically acceptable salts thereof:

15



(I)

wherein A is a phenyl or heterocyclic group, optionally substituted with 1 to 4  
20 substituents selected from the group consisting of a halogen atom, a hydroxyl group, an amino group, a nitro group, a cyano group, an alkyl group having 1 to 4 carbons, an alkoxy group having 1 to 4 carbons, an aminoalkyl group having 1 to 4 carbons, an alkylamino group having 1 to 4 carbons, an acyl group having 2 to 4 carbons, an acylamino group having 2 to 4 carbons, an alkythio group having 1 to 4 carbons, a

perfluoroalkyl group having 1 to 4 carbons, a perfluoroalkyloxy group having 1 to 4 carbons, a carboxyl group, an alkoxycarbonyl group having 1 to 4 carbons, a phenyl group and a heterocyclic group;

B is a phenyl or heterocyclic group, optionally substituted with 1 to 3 substituents  
5 selected from the group consisting of a halogen atom, a hydroxyl group, an amino group, a nitro group, a cyano group, an alkyl group having 1 to 4 carbons, an alkoxy group having 1 to 4 carbons, an aminoalkyl group having 1 to 4 carbons, an alkylamino group having 1 to 4 carbons, an acyl group having 2 to 4 carbons, an acylamino group having 2 to 4 carbons, an alkythio group having 1 to 4 carbons, a perfluoroalkyl group  
10 having 1 to 4 carbons, a perfluoroalkyloxy group having 1 to 4 carbons, a carboxyl group, an alkoxycarbonyl group having 1 to 4 carbons, a phenyl group and a heterocyclic group;

Z is a bond, an optionally substituted alkylene having 1 to 4 carbons or a moiety having -O-, -S-, -NH-, -CO-, -CS-, -SO- or -SO<sub>2</sub>- which is linear, cyclic or their  
15 combination;

Y is a moiety having -CO-, -CS-, -SO- or -SO<sub>2</sub>- which is linear, cyclic or their combination; and in which the distances between the centroid of ring B (W1), the centroid of ring A (W2) and an oxygen or sulfur atom as a hydrogen bond acceptor in the moiety Y (W3) can, for example, be as follows: W1-W2=6.0 to 12.0 Å,  
20 W1-W3=3.0 to 6.0 Å, and W2-W3=4.0 to 8.0 Å; preferably W1-W2=8.0 to 10.0 Å, W1-W3=3.0 to 5.0 Å, W2-W3=5.0 to 8.0 Å (compounds of the invention described herein, however, are not necessarily limited to these dimensions);

R<sup>1</sup> and R<sup>2</sup> are independently a hydrogen or an optionally substituted alkyl having 1 to 4 carbons; or R<sup>1</sup> and R<sup>2</sup> may form a bond;

25 R<sup>3</sup> is a hydrogen or an optionally substituted alkyl having 1 to 4 carbons; R<sup>4</sup> is a hydrogen atom, halogen atom, a hydroxyl group, an amino group, a nitro group, a

cyano group, an alkyl group having 1 to 4 carbons, an alkoxy group having 1 to 4 carbons, an aminoalkyl group having 1 to 4 carbons, an alkylamino group having 1 to 4 carbons, an acyl group having 1 to 4 carbons, an acylamino group having 1 to 4 carbons, an alkylthio group having 1 to 4 carbons, a perfluoroalkyl group having 1 to 4 carbons, a perfluoroalkyloxy group having 1 to 4 carbons, a carboxyl group or an alkoxycarbonyl group having 1 to 4 carbons.

One of  $X^1$ ,  $X^2$ ,  $X^3$ , or  $X^4$  is a halogen atom, a hydroxyl group, an amino group, a nitro group, a cyano group, an alkyl group having 1 to 4 carbons, an alkoxy group having 1 to 4 carbons, an aminoalkyl group having 1 to 4 carbons, an alkylamino group having 1 to 4 carbons, an acyl group having 1 to 4 carbons, an acylamino group having 1 to 4 carbons, an alkylthio group having 1 to 4 carbons, a perfluoroalkyl group having 1 to 4 carbons, a perfluoroalkyloxy group having 1 to 4 carbons, a carboxyl group or an alkoxycarbonyl group having 1 to 4 carbons, while the others of  $X^1$ ,  $X^2$ ,  $X^3$ , or  $X^4$  independently are a hydrogen atom, a halogen atom, a hydroxyl group, an amino group, a nitro group, a cyano group, an alkyl group having 1 to 4 carbons, an alkoxy group having 1 to 4 carbons, an aminoalkyl group having 1 to 4 carbons, an alkylamino group having 1 to 4 carbons, an acyl group having 1 to 4 carbons, an acylamino group having 1 to 4 carbons, an alkylthio group having 1 to 4 carbons, a perfluoroalkyl group having 1 to 4 carbons, a perfluoroalkyloxy group having 1 to 4 carbons, a carboxyl group or an alkoxycarbonyl group having 1 to 4 carbons.

In the above structural formula (I) and throughout the present specification, the following terms have the indicated meaning:

The term "heterocyclyl" as used herein means a monovalent saturated or unsaturated group being monocyclic and containing one or more heteroatoms, such as pyrrolidine, pyrroline, pyrazoline, imidazolidine, imidazoline, piperidine, morpholine and the like.

The term "halogen" as used herein means fluorine, chlorine, bromine or iodine.

The term "alkyl having 1 to 4 carbons" as used herein includes methyl, ethyl, n-propyl, iso-propyl, butyl, iso-butyl, sec-butyl and tert-butyl.

The term "alkoxy having 1 to 4 carbons" as used herein includes methoxy, ethoxy,  
5 n-propoxy, isopropoxy, n-butoxy, isobutoxy and the like.

The term "aminoalkyl having 1 to 4 carbons" as used herein includes aminomethyl, 1-aminopropyl, 2-aminopropyl and the like.

The term "alkylamino having 1 to 4 carbons" as used herein includes N-methylamino, N-ethylamino, N-isopropylamino and the like.

10 The term "acyl having 2 to 4 carbons" as used herein includes acetyl, propionyl, butyryl, isobutyryl and the like.

The term "acylamino having 2 to 4 carbons" as used herein includes acetalamino, propionylamino, butyrylamino, isobutyrylamino and the like.

The term "alkylthio having 2 to 4 carbons" as used herein includes methylthio,  
15 ethylthio, propylthio and the like.

The term "perfluoroalkyl having 2 to 4 carbons" as used herein includes trifluoromethyl, pentafluoroethyl and the like.

The term "perfluoroalkyloxy having 2 to 4 carbons" as used herein includes trifluoromethoxy, pentafluoroethoxy and the like.


20 The term "alkylene having 1 to 4 carbons" as used herein includes methylene, ethylene and the like.

The term "ring centroid" used in definition of the spatial configuration may be defined as an average of X, Y and Z axes of the ring-forming atoms.



25 The compounds of this invention are prepared as described below:

(a) A compound represented by formula (II) is condensed with a compound

represented by formula (III) to give a compound represented by formula (IV):



5




10

wherein A, Z, Y, B, R<sup>1</sup> and R<sup>2</sup> are as defined above; R<sup>5</sup> is a moiety having -C(=Q)OH (Q is an oxygen or sulfur atom) or a moiety having -NH<sub>2</sub>; R<sup>6</sup> is a moiety having -NH<sub>2</sub> when R<sup>5</sup> is a moiety having -C(=Q)OH (Q is an oxygen or sulfur atom) and a moiety having -C(=Q)OH (Q is an oxygen or sulfur atom) when R<sup>5</sup> is a moiety having -NH<sub>2</sub>.

15

(b) A compound represented by formula (IV) is condensed with a compound represented by formula (V) to give the compound of this invention.



20

wherein R<sup>3</sup>, R<sup>4</sup>, X<sup>1</sup>, X<sup>2</sup>, X<sup>3</sup>, and X<sup>4</sup> are as defined above.

The above condensation reactions (a) and (b) are conducted using a peptide condensing agent such as dicyclohexylcarbodiimide, N,N'-carbonyldiimidazole, diphenyl

phosphoric azide, diethylphosphorylcyanide, etc.

The reaction may be conducted at 0 to 80 °C for 4 to 72 hours. Solvents which may be used are normal solvents such as benzene, toluene, tetrahydrofuran, dioxane,  
5 dichloromethane, chloroform, N, N-dimethylformamide, etc. If necessary, a base such as sodium hydroxide, triethylamine and pyridine or an acid such as hydrochloric acid, acetic acid and trifluoroacetic acid may be added to the reaction system.

The compound of this invention and the intermediate represented by formula (I) may  
10 be purified or isolated by the conventional separation method such as extraction, recrystallization, column chromatography and the like.

The novel compounds of this invention have differentiation-inducing effects and thus are useful as therapeutic treatment or improving agents related to differentiation and/or  
15 proliferation-related disorders such as cancer and psoriasis. In particular, they are highly effective as carcinostatic agents to hematological malignancy and solid carcinoma.


The active ingredient of this invention useful as a drug may be used in the form of a  
20 general pharmaceutical composition. The pharmaceutical composition may be in the forms normally employed, such as tablets, capsules, powders, syrups, solutions, suspensions, aerosols, and the like, may contain flavourants, sweeteners, etc. in suitable solids or liquid carriers or diluents, or in suitable sterile media to form injectable solutions or suspensions. Such composition typically contains from 1 to 70%,  
25 preferably 5 to 50% by weight of active compound, the remainder of the composition being pharmaceutically acceptable carriers, diluents or solvents or salt solutions.

The compounds of this invention are clinically administered to mammals, including

man and animals, via oral, nasal, transdermal, pulmonary, or parenteral routes. Administration by the oral route is preferred, being more convenient and avoiding the possible pain and irritation of injection. By either route, the dosage is in the range of about 0.0001 to about 200 mg/kg body weight per day administered singly or as a  
5 divided dose. However, the optimal dosage for the individual subject being treated will be determined by the person responsible for treatment, generally smaller doses being administered initially and thereafter increments made to determine the most suitable dosage.

10 The following examples are given as specific illustrations of the invention. It should be understood, however, that the invention is not limited to the specific details set forth in the examples. All parts and percentages in the examples, as well as in the remainder of the specification, are by weight unless otherwise specified.

15 Further, any range of numbers recited in the specification or paragraphs hereinafter describing or claiming various aspects of the invention, such as that representing a particular set of properties, units of measure, conditions, physical states or percentages, is intended to literally incorporate expressly herein by reference or otherwise, any number falling within such range, including any subset of numbers or ranges subsumed  
20 within any range so recited. The term "about" when used as a modifier for, or in conjunction with, a variable, is intended to convey that the numbers and ranges disclosed herein are flexible and that practice of the present invention by those skilled in the art using temperatures, concentrations, amounts, contents, carbon numbers, and properties that are outside of the range or different from a single value, will achieve the  
25 desired result.


**Example 1.****Preparation of 4-[N-(Pyridin-3-ylacryloyl)aminomethyl]benzoic acid**

5

To a suspension of 0.33 g (2.01 mmol) of N, N'-carbonyldiimidazole in tetrahydrofuran (10 ml) is added drop-wise a solution of 0.30 g (2.01 mmol) of 3-pyridineacrylic acid at 0 °C. Then, the mixture is stirred at room temperature for 3 hours and added drop-wise to a separately prepared 2.0 ml (2.00 mmol) of 1N aqueous sodium hydroxide solution including 0.30 g (2.00 mmol) of 4-aminomethylbenzoic acid, followed by stirring at room temperature for 8 hours. The reaction mixture is evaporated under vacuum. To the residue is added a saturated solution of sodium chloride (2 ml), then the mixture is neutralized with concentrated hydrochloric acid to pH 5. The deposited white solid is collected by filtration, washed with ice-water, and then dried to give the title compound (0.46 g, 82%). HRMS calcd for C<sub>16</sub>H<sub>14</sub>N<sub>2</sub>O<sub>3</sub>: 282.2988. Found: 282.2990. MA calcd for: C<sub>16</sub>H<sub>14</sub>N<sub>2</sub>O<sub>3</sub>: C, 68.07%; H, 5.00%; N, 9.92. Found: C, 68.21%; H, 5.03%; N, 9.90%.

10

15

**Example 2. Preparation of**20 **N-(2-amino-5-fluorophenyl)-4-[N-(Pyridn-3-ylacryloyl)aminomethyl]benzamide**




To a suspension of 0.29 g (1.78 mmol) of N, N'-carbonyldiimidazole in tetrahydrofuran (15 ml) is added 0.50 g (1.78 mmol) of 4-[N-(Pyridn-3-ylacryloyl)aminomethyl]benzoic acid, followed by stirring at 45°C for 1 hour. After cooling, the reaction mixture is added to a separately prepared  
5 tetrahydrofuran (10 ml) solution including 0.28 g (2.22 mmol) of 4-fluoro-1,2-phenylenediamine and 0.20 g (1.78 mmol) of trifluoroacetic acid at room temperature. After reaction at room temperature for 24 hours, the deposited white solid is collected by filtration, washed with tetrahydrofuran, and then dried to give the title compound (0.40 g, 57%). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δppm: 4.49 (2H, d), 4.84  
10 (2H, br.s), 6.60 (1H, t), 6.80 (2H, m), 6.96 (1H, t), 7.18 (1H, d), 7.42 (2H, d), 7.52 (1H, d), 7.95 (2H, d), 8.02 (1H, d), 8.56 (1H, d), 8.72 (1H, br. t), 8.78 (1H, s), 9.60 (1H, br.s). IR (KBr) cm<sup>-1</sup>: 3310, 1655, 1631, 1524, 1305, 750. HRMS calcd for C<sub>22</sub>H<sub>19</sub>N<sub>4</sub>O<sub>2</sub>F: 390.4170. Found: 390.4172. MA calcd for C<sub>22</sub>H<sub>19</sub>N<sub>4</sub>O<sub>2</sub>F: C, 67.68%; H, 4.40%; N, 14.35. Found: C, 67.52%; H, 4.38%; N, 14.42%.

15

**Example 3.**

Preparation of 4-[N-cinnamoylaminomethyl]benzoic acid




20 To a suspension of 0.33 g (2.01 mmol) of N, N'-carbonyldiimidazole in tetrahydrofuran (10 ml) is added drop-wise a solution of 0.30 g (2.01 mmol) of cinnamic acid at 0 °C. Then, the mixture is stirred at room temperature for 3 hours and added drop-wise to a separately prepared 2.0 ml (2.00 mmol) of 1N aqueous sodium hydroxide solution including 0.30 g (2.00 mmol) of 4-aminomethylbenzoic acid,  
25 followed by stirring at room temperature for 8 hours. The reaction mixture is

evaporated under vacuum. To the residue is added a saturated solution of sodium chloride (2 ml), then the mixture is neutralized with concentrated hydrochloric acid to pH 7. The deposited white solid is collected by filtration, washed with ice-water, and then dried to give the title compound (0.51 g, 91%). HRMS calcd for  $C_{17}H_{15}NO_3$ : 281.3242. Found: 281.3240. MA calcd for  $C_{17}H_{15}NO_3$ : C, 72.58%; H, 5.38%; N, 4.98. Found: C, 72.42%; H, 5.37%; N, 4.87%.

#### Example 4.

Preparation of N-(2-amino-5-fluorophenyl)-4-[N-cinnamoylaminomethyl]benzamide



To a suspension of 0.29 g (1.78 mmol) of N, N'-carbonyldiimidazole in tetrahydrofuran (15 ml) is added 0.50 g (1.78 mmol) of 4-[N-cinnamoylaminomethyl]benzoic acid, followed by stirring at 45°C for 1 hour. After cooling, the reaction mixture is added to a separately prepared tetrahydrofuran (10 ml) solution including 0.28 g (2.22 mmol) of 4-fluoro-1,2-phenylenediamine and 0.20 g (1.78 mmol) of trifluoroacetic acid at room temperature. After reaction at room temperature for 16 hours, the deposited white solid is collected by filtration, washed with tetrahydrofuran, and then dried to give the title compound (0.45 g, 64%).  $^1H$  NMR (300 MHz, DMSO- $d_6$ ):  $\delta$ ppm: 4.42 (2H, d), 4.92 (2H, br.s), 6.62 (1H, t), 6.78 (2H, m), 7.01 (1H, t), 7.32 (5H, m), 7.54 (5H, m), 8.76 (1H, br.t), 9.58 (1H, br.s). IR (KBr)  $cm^{-1}$ : 3306, 1618, 1517, 1308, 745. HRMS calcd for  $C_{23}H_{20}N_3O_2F$ : 389.4292. Found: 389.4294. MA calcd for  $C_{23}H_{20}N_3O_2F$ : C, 70.94%; H, 5.18%; N, 10.79. Found: C,

70.72%; H, 5.18%; N, 10.88%.

### Example 5.

In vitro inhibition of HDAC enzymatic activity by  
 5 N-(2-amino-5-fluorophenyl)-4-[N-(Pyridin-3-ylacryloyl)aminomethyl] benzamide  
 (Compound CS02100055),  
 N-(2-aminophenyl)-4-[N-(4-fluorophenyl)aminomethyl]benzamide (compound  
 CS02100019), and N-(2-aminophenyl)-4-[N-(pyridin-3-ylmethoxycarbonyl)  
 aminomethyl]benzamide (MS-275, EP0847992).

10

The inhibitory effects of MS-275 and compounds CS02100055 and CS02100019 on  
 HDAC were tested by a HDAC colorimetric activity assay kit (BIOMOL Research  
 Laboratories, PA, USA) according to the instructions from the manufacturer. Briefly,  
 15 the tested compounds at different concentrations were added to 96-well plates, then  
 mixed with extract from HeLa cells containing HDAC activity provided by the  
 manufacturer. HDAC reactions were initiated by adding substrate. 10 minutes later, the  
 reactions were stopped by addition of Color De Lys Developer. Microplates were read  
 in a plate reader at 405 nm. Inhibition of HDAC activity was calculated following the  
 20 instructions. The testing results are listed in Table 1.

**Table 1. *In vitro* Inhibition of HDAC Activity by MS275, CS02100055 and CS02100019**

Compound	% of inhibition of HDAC activity at different concentrations				IC <sub>50</sub> ( $\mu$ M)
	1 $\mu$ M	5 $\mu$ M	10 $\mu$ M	50 $\mu$ M	
MS-275	14.7	19.1	64.1	82.3	8.4
CS02100055	17.5	37.3	62.1	78.4	7.2
CS02100019	11.3	24.9	28.4	29.8	>50.0

**Example 6.**

The growth inhibitory effect of N-(2-amino-5-fluorophenyl)-4-

[N-(Pyridin-3-ylacryloyl) aminomethyl] benzamide (Compound CS02100055),

- 5 N-(2-aminophenyl)-4-[N-(4-fluorophenyl)aminomethyl]benzamide (Compound CS02100019), and N-(2-aminophenyl)-4-[N-(pyridin-3-ylmethoxycarbonyl) aminomethyl]benzamide (MS-275, EP0847992), on various tumor cell lines *in vitro*.

Growth inhibition tests were carried out by MTS method. Approximately 72 hours  
10 before the viability assay, cells were seeded into 96-well plates at  $5-10 \times 10^3$  cells/well (according to the growth rate of individual cell lines used). 24 hours later, the tested compounds at different concentrations were added, and the cells were cultured for 48 hours, then added 20  $\mu$ l/well of CellTiter 96 AQueous One Solution Reagent containing tetrazolium compound (Promega) into each well. MTS was subsequently added to the  
15 culture medium. After incubation of the plates for 2 hours at about 37°C, the absorbance at 490 nm was recorded by a 96-well plate reader. Cell viability was calculated by  $A_{\text{treatment}}/A_{\text{control}} \times 100\%$  ( $A$  represents the absorbance recorded at 490 nm). The concentration that inhibited cell growth by 50% over the control was determined as  $GI_{50}$ . All of the compounds were dissolved in DMSO and were added to the culture at a  
20 1:1000 dilution to give a final DMSO concentration of  $\leq 0.1\%$ . All samples were assayed in duplicate, and each experiment was repeated at least three times. The testing results are summarized in Table 2.

**Table 2. Growth Inhibition of Tumor Cell lines by MS-275, CS2100055 and CS02100019**

Compound	GI <sub>50</sub> (μM) against Various Tumor Cell Lines*							
	U2OS	HeLa	DU-145	SMMC-7721	HepG2	293	MCF-7	231
MS-275	1.0	25	13	20	3.2	16	6.3	5.0
CS02100055	2.0	40	25	16	4.0	50	5.0	7.9
CS02100019	2.5	50	50	50	3.2	25	5.0	7.9

Compound	GI <sub>50</sub> (μM) against Various Tumor Cell Lines*							
	LNCaP	SK-N-SH	PANC-1	SK-OV-3	SGC-7901	Raji	HL-60	28SC
MS-275	2.5	50	5.0	50	50	6.3	0.32	4.0
CS02100055	4.0	50	6.3	50	50	4.0	0.4	5.8
CS02100019	10	50	5.0	50	50	2.0	0.5	2.0

\* Origin of the cell lines:

U2OS, human osteocarcinoma

HeLa, human cervical carcinoma

5 DU-145, human prostate cancer

SGC-7901, human gastric adenocarcinoma

SMMC-7721, human hepatoma

HepG2, human hepatoblastoma

293, human embryonic kidney

MCF-7, human breast adenocarcinoma

MDA-MB-231, human breast adenocarcinoma H292, human lung cancer

LNCaP, human prostate cancer

SK-N-SH, human neuroblastoma

10 PANC-1, human pancreatic ductal carcinoma

SK-OV-3, human ovary adenocarcinoma

28SC, human macrophage

Raji, human Burkitts Lymphoma

HL-60, human myeloid leukemia

Jurkat, human T cell leukemia

### Example 7.

15 Transcription activation of nuclear hormone receptors by  
N-(2-amino-5-fluorophenyl)-4- [N-(Pyridin-3-ylacryloyl) aminomethyl] benzamide  
(Compound CS02100055), N-(2-aminophenyl)-4-[N-(pyridin-3-ylmethoxycarbonyl)  
aminomethyl]benzamide (MS-275, EP0847992) and Trichostatin A (TSA).

20 Transcription activation of several nuclear hormone receptors by tested


compounds, as indicated in Figure 1, was carried out by reporter assay experiments. Briefly, U2OS cells were seeded in 96-well plates the day before transfection to give a confluency of 50-80%. Cells were transfected with one of the expression plasmids containing cDNAs coding either glucocorticoid receptor (GR), peroxisome proliferator activated receptor  $\gamma$  (PPAR $\gamma$ , estrogen receptor  $\alpha$  (ER $\alpha$ ), or estrogen receptor  $\beta$  (ER  $\beta$ ), in combination with retinoid X receptor  $\alpha$  (RXR  $\alpha$ ), and their corresponding luciferase reporter plasmids using FuGene6 transfection reagent according to the manufacturers instructions (Roche). Cells were allowed to express protein for 24 hours followed by addition of individual compounds or the vehicle (DMSO). 24 hours later cells were harvested, and the luciferase assays were performed using the luciferase assay kit according to the manufacturer's instructions (Promega). To normalize the data from the luciferase assays,  $\beta$ -galactosidase activity from transfected cells was measured using a kit (Promega) as instructed by the manufacturer. Response elements for individual nuclear receptors were as following: GR (5'-GATCTTGTACAGGATGTTCTCTAGCGATGTACAGGATGTTCTCTAGCGATGTACAGGATGTTCTCTAG-3') (SEQ ID NO. 1), PPAR (5'-CGCGTTCCTTTCCGAACGTGACCTTTGTCCTGGTCCCCTTTTGCT-3') (SEQ ID NO. 2), and ER (5'-TCGAGTCAGGTCACAGTGACCTGATC-3') (SEQ ID NO. 3). The testing results are summarized in Figure 1. FIG.1 shows transcription activation of nuclear hormone receptors by different HDAC inhibitors Trichostatin A, MS-275 and CS2100055. Experiments were carried out as described above. LD stands for corresponding Ligands for each receptor, and CS55 for CS2100055 in each panel of the figure. Concentrations of tested compounds in all experiments were TSA 0.2  $\mu$ M, MS-275 1  $\mu$ M, and CS55 1  $\mu$ M. Dexamethasone (0.1  $\mu$ M), Rosiglitazone (10  $\mu$ M), and E2 (0.01  $\mu$ M) were used as ligands for GR, PPAR $\gamma$ , and ER, respectively. Three independent experiments were performed and the results from a representative

experiment were shown in Figure 1.

\* \* \*

What is claimed is:

1. A compound of formula I:



(I)

or its stereoisomer, enantiomer, diastereomer, hydrate, or a pharmaceutically acceptable salt;

10 wherein A is a phenyl or heterocyclic group, optionally substituted with 1 to 4 substituents selected from the group consisting of a halogen atom, a hydroxyl group, an amino group, a nitro group, a cyano group, an alkyl group having 1 to 4 carbons, an alkoxy group having 1 to 4 carbons, an aminoalkyl group having 1 to 4 carbons, an alkylamino group having 1 to 4 carbons, an acyl group having 2 to 4 carbons, an acylamino group having 2 to 4 carbons, an alkythio group having 1 to 4 carbons, a perfluoroalkyl group having 1 to 4 carbons, a perfluoroalkyloxy group having 1 to 4 carbons, a carboxyl group, an alkoxycarbonyl group having 1 to 4 carbons, a phenyl group and a heterocyclic group;

20 B is a phenyl or heterocyclic group, optionally substituted with 1 to 3 substituents selected from the group consisting of a halogen atom, a hydroxyl group, an amino group, a nitro group, a cyano group, an alkyl group having 1 to 4 carbons, an alkoxy group having 1 to 4 carbons, an aminoalkyl group having 1 to 4 carbons, an alkylamino group having 1 to 4 carbons, an acyl group having 2 to 4 carbons, an acylamino group having 2 to 4 carbons, an alkythio group having 1 to 4 carbons, a perfluoroalkyl group



having 1 to 4 carbons, a perfluoroalkyloxy group having 1 to 4 carbons, a carboxyl group, an alkoxycarbonyl group having 1 to 4 carbons, a phenyl group and a heterocyclic group;

5        Z is a bond, an optionally substituted alkylene having 1 to 4 carbons or a moiety having -O-, -S-, -NH-, -CO-, -CS-, -SO- or -SO<sub>2</sub>- which is linear, cyclic or their combination;

Y is a moiety having -CO-, -CS-, -SO- or -SO<sub>2</sub>- which is linear, cyclic or a combination thereof; and in which the distances between the centroid of ring B (W1),  
10       the centroid of ring A (W2) and an oxygen or sulfur atom as a hydrogen bond acceptor in the moiety Y (W3) are: W1-W2= about 6.0 to about 12.0 Å, W1-W3= about 3.0 to about 6.0 Å, and W2-W3= about 4.0 to about 8.0 Å, respectively; or W1-W2= about 8.0 to about 10.0 Å, W1-W3= about 3.0 to about 5.0 Å, W2-W3= about 5.0 to about 8.0 Å, respectively;

15       R<sup>1</sup> and R<sup>2</sup> are independently a hydrogen or an optionally substituted alkyl having 1 to 4 carbons; or R<sup>1</sup> and R<sup>2</sup> may form a bond;


R<sup>3</sup> is a hydrogen or an optionally substituted alkyl having 1 to 4 carbons;

R<sup>4</sup> is a hydrogen atom, halogen atom, a hydroxyl group, an amino group, a nitro group, a cyano group, an alkyl group having 1 to 4 carbons, an alkoxy group having 1  
20       to 4 carbons, an aminoalkyl group having 1 to 4 carbons, an alkylamino group having 1 to 4 carbons, an acyl group having 1 to 4 carbons, an acylamino group having 1 to 4 carbons, an alkylthio group having 1 to 4 carbons, a perfluoroalkyl group having 1 to 4 carbons, a perfluoroalkyloxy group having 1 to 4 carbons, a carboxyl group or an alkoxycarbonyl group having 1 to 4 carbons; and

25       One of X<sup>1</sup>, X<sup>2</sup>, X<sup>3</sup>, or X<sup>4</sup> is a halogen atom, a hydroxyl group, an amino group, a nitro group, a cyano group, an alkyl group having 1 to 4 carbons, an alkoxy group

having 1 to 4 carbons, an aminoalkyl group having 1 to 4 carbons, an alkylamino group having 1 to 4 carbons, an acyl group having 1 to 4 carbons, an acylamino group having 1 to 4 carbons, an alkylthio group having 1 to 4 carbons, a perfluoroalkyl group having 1 to 4 carbons, a perfluoroalkyloxy group having 1 to 4 carbons, a carboxyl group or an alkoxycarbonyl group having 1 to 4 carbons, while the others of X<sup>1</sup>, X<sup>2</sup>, X<sup>3</sup>, or X<sup>4</sup> independently are a hydrogen atom, a halogen atom, a hydroxyl group, an amino group, a nitro group, a cyano group, an alkyl group having 1 to 4 carbons, an alkoxy group having 1 to 4 carbons, an aminoalkyl group having 1 to 4 carbons, an alkylamino group having 1 to 4 carbons, an acyl group having 1 to 4 carbons, an acylamino group having 1 to 4 carbons, an alkylthio group having 1 to 4 carbons, a perfluoroalkyl group having 1 to 4 carbons, a perfluoroalkyloxy group having 1 to 4 carbons, a carboxyl group or an alkoxycarbonyl group having 1 to 4 carbons.

2. A compound of formula IV:



or its stereoisomer, enantiomer, diastereomer, hydrate, or a pharmaceutically acceptable salt;

wherein A is a phenyl or heterocyclic group, optionally substituted with 1 to 4 substituents selected from the group consisting of a halogen atom, a hydroxyl group, an amino group, a nitro group, a cyano group, an alkyl group having 1 to 4 carbons, an alkoxy group having 1 to 4 carbons, an aminoalkyl group having 1 to 4 carbons, an alkylamino group having 1 to 4 carbons, an acyl group having 2 to 4 carbons, an acylamino group having 2 to 4 carbons, an alkylthio group having 1 to 4 carbons, a perfluoroalkyl group having 1 to 4 carbons, a perfluoroalkyloxy group having 1 to 4

carbons, a carboxyl group, an alkoxycarbonyl group having 1 to 4 carbons, a phenyl group and a heterocyclic group;

B is a phenyl or heterocyclic group, optionally substituted with 1 to 3 substituents selected from the group consisting of a halogen atom, a hydroxyl group, an amino  
5 group, a nitro group, a cyano group, an alkyl group having 1 to 4 carbons, an alkoxy group having 1 to 4 carbons, an aminoalkyl group having 1 to 4 carbons, an alkylamino group having 1 to 4 carbons, an acyl group having 2 to 4 carbons, an acylamino group having 2 to 4 carbons, an alkythio group having 1 to 4 carbons, a perfluoroalkyl group having 1 to 4 carbons, a perfluoroalkyloxy group having 1 to 4 carbons, a carboxyl  
10 group, an alkoxycarbonyl group having 1 to 4 carbons, a phenyl group and a heterocyclic group;




Z is a bond, an optionally substituted alkylene having 1 to 4 carbons or a moiety having -O-, -S-, -NH-, -CO-, -CS-, -SO- or -SO<sub>2</sub>- which is linear, cyclic or their combination;

15 Y is a moiety having -CO-, -CS-, -SO- or -SO<sub>2</sub>- which is linear, cyclic or a combination thereof; and in which the distances between the centroid of ring B (W1), the centroid of ring A (W2) and an oxygen or sulfur atom as a hydrogen bond acceptor in the moiety Y (W3) is: W1-W2= about 6.0 to about 12.0 Å, W1-W3= about 3.0 to about 6.0 Å, and W2-W3= about 4.0 to about 8.0 Å, respectively; or W1-W2= about  
20 8.0 to about 10.0 Å, W1-W3= about 3.0 to about 5.0 Å, W2-W3= about 5.0 to about 8.0 Å, respectively; and

R<sup>1</sup> and R<sup>2</sup> are independently a hydrogen or an optionally substituted alkyl having 1 to 4 carbons; Or R<sup>1</sup> and R<sup>2</sup> may form a bond.


25 3. A process for the preparation of a compound according to claim 1 or its stereoisomer, enantiomer, diastereomer, hydrate or pharmaceutically acceptable salt thereof comprising the steps of:

- (a) condensing a compound represented by formula (II) with a compound represented by formula (III) to yield a compound represented by formula (IV):



$\text{R}^5$  is a moiety having  $-\text{C}(=\text{Q})\text{OH}$  wherein Q is an oxygen or sulfur atom, or  $\text{R}^5$  is a moiety having  $-\text{NH}_2$ ;  $\text{R}^6$  is a moiety having  $-\text{NH}_2$  when  $\text{R}^5$  is a moiety having  $-\text{C}(=\text{Q})\text{OH}$  wherein Q is an oxygen or sulfur atom, or a moiety having  $-\text{C}(=\text{Q})\text{OH}$  wherein Q is an oxygen or sulfur atom when  $\text{R}^5$  is a moiety having  $-\text{NH}_2$ ; and

- (b) condensing the compound represented by formula (IV) with a compound represented by formula (V) to give the compound of formula (I).



wherein one of  $\text{X}^1$ ,  $\text{X}^2$ ,  $\text{X}^3$ , or  $\text{X}^4$  is a halogen atom, a hydroxyl group, an amino

group, a nitro group, a cyano group, an alkyl group having 1 to 4 carbons, an alkoxy group having 1 to 4 carbons, an aminoalkyl group having 1 to 4 carbons, an alkylamino group having 1 to 4 carbons, an acyl group having 1 to 4 carbons, an acylamino group having 1 to 4 carbons, an alkylthio group having 1 to 4 carbons, a perfluoroalkyl group having 1 to 4 carbons, a perfluoroalkyloxy group having 1 to 4 carbons, a carboxyl group or an alkoxycarbonyl group having 1 to 4 carbons, and the other of  $X^1$ ,  $X^2$ ,  $X^3$ , or  $X^4$  independently are a hydrogen atom, a halogen atom, a hydroxyl group, an amino group, a nitro group, a cyano group, an alkyl group having 1 to 4 carbons, an alkoxy group having 1 to 4 carbons, an aminoalkyl group having 1 to 4 carbons, an alkylamino group having 1 to 4 carbons, an acyl group having 1 to 4 carbons, an acylamino group having 1 to 4 carbons, an alkylthio group having 1 to 4 carbons, a perfluoroalkyl group having 1 to 4 carbons, a perfluoroalkyloxy group having 1 to 4 carbons, a carboxyl group or an alkoxycarbonyl group having 1 to 4 carbons;

$R^3$  is a hydrogen or an optionally substituted alkyl having 1 to 4 carbons; and

$R^4$  is a hydrogen atom, halogen atom, a hydroxyl group, an amino group, a nitro group, a cyano group, an alkyl group having 1 to 4 carbons, an alkoxy group having 1 to 4 carbons, an aminoalkyl group having 1 to 4 carbons, an alkylamino group having 1 to 4 carbons, an acyl group having 1 to 4 carbons, an acylamino group having 1 to 4 carbons, an alkylthio group having 1 to 4 carbons, a perfluoroalkyl group having 1 to 4 carbons, a perfluoroalkyloxy group having 1 to 4 carbons, a carboxyl group or an alkoxycarbonyl group having 1 to 4 carbons.

4. The process according to claim 3, wherein the condensation reactions of steps (a) and (b) are carried out using a peptide condensing agent.

25

5. The process according to claim 4, wherein said peptide condensing agent is dicyclohexylcarbodiimide, N, N'-carbonyldiimidazole, diphenyl phosphoric azide, or

diethylphosphorylcyanide.

6. The process of claim 3, wherein said condensation reactions of steps (a) and (b) are conducted at a temperature from about 0°C to about 80°C.

5

7. A pharmaceutical composition useful as a therapeutic and/or improving agent for differentiation and/or proliferation-related disorders comprising an effective amount of a compound according to claim 1 and at least one pharmaceutically acceptable excipient, carrier or diluent.

10

8. The pharmaceutical composition of claim 7, wherein said proliferation-related disorder is selected from the group consisting essentially of psoriasis, a hematological malignancy and a solid carcinoma.

15 9. A dosage form unit of the pharmaceutical composition of claim 7 comprising an amount within the range of about 0.0001 to about 200 mg of said compound.

10. A pharmaceutical composition according to claim 7 for administration by the oral, nasal, transdermal, pulmonary, or parenteral route.

20

11. A method of treating cell proliferative diseases comprising administering to a subject in need thereof an effective amount of a compound of claim 1.

12. A method according to claim 11, wherein said cell proliferative disease is selected  
25 from the group consisting essentially of a malignant tumor and psoriasis.

13. The method according to claim 11, wherein the effective amount of the compound is within the range of about 0.0001 to about 200 mg/kg body weight per day.

14. A method of treating a cell proliferative disease comprising administering to a subject in need thereof an effective amount of a compound according to claim 1 in combination with an active pharmaceutical chemotherapeutic compound or methyl transferase inhibitor formulated with at least one pharmaceutically acceptable excipient, carrier or diluent.
15. The method of claim 14, wherein said cell proliferative disease is selected from the group consisting essentially of a malignant tumor and psoriasis.
16. A pharmaceutical composition for activating nuclear receptors comprising an effective amount of a compound according to claim 1 and at least one a pharmaceutically acceptable excipient, carrier or diluent.
17. A dosage form unit of the pharmaceutical composition of claim 16 comprising an amount within the range of about 0.0001 to about 200 mg of the compound.
18. A pharmaceutical composition according to claim 16 for administration by the oral, nasal, transdermal, pulmonary, or parenteral route.
19. A method of treatment or prevention of a condition mediated by a nuclear receptor comprising administering to a subject in need thereof an effective amount of a compound of claim 1.
20. A method of treatment or prevention of a condition mediated by abnormally low activity of a nuclear receptor comprising administering to a subject in need thereof an effective amount of a compound according to claim 1.


21. A method according to claim 19, wherein the condition is selected from the group consisting essentially of an endocrine-related disorder, an immune system or inflammatory disorder, a genetic disorder, and neurodegeneration.
- 5 22. A method according to claim 20, wherein the condition is selected from the group consisting essentially of an endocrine-related disorder, an immune system or inflammatory disorder, a genetic disorder, and neurodegeneration.
23. The method according to claim 19, wherein the effective amount of the compound  
10 is within the range of about 0.0001 to about 200mg/kg body weight per day.
24. The method according to claim 20, wherein the effective amount of the compound is within the range of about 0.0001 to about 200mg/kg body weight per day

15

\* \* \*



FIG.1



# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/IB2004/000401

## A. CLASSIFICATION OF SUBJECT MATTER

IPC7 C07D213/56, C07C233/51, C07C237/40, A61K31/4406, 31/165, A61P17/06, 35/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7 C07C C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

STN CA plus

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 01 16106A1 (SCHERING AG), 08.Mar 2001(08.03.2001) see the whole document	1-24
A	EP 0847992 A1 (SCHERING AG) 17.Jun 1998(17.06.1998) see the whole document	1-24

☐ Further documents are listed in the continuation of Box C. ☒ See patent family annex.

### \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim (S) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

28 Mar 2005 (28.03.2005)

Date of mailing of the international search report

21 · APR 2005 (21 · 04 · 2005)

Name and mailing address of the ISA/CN

6 Xitucheng Rd., Jimen Bridge, Haidian District,  
100088 Beijing, China  
Facsimile No. 86-10-62019451

Authorized officer

Hou Yao

Telephone No. 86-10-62085608

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/IB2004/000401

## Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 1,2  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:  
The definitions of moieties "Z" and "Y" of general formulas in claims 1 and 2 are vague and unclear that skilled in the art could not figure out the exact structure of the moieties, thereby render the definition of the subject-matter of said claims unclear. This report is on the basis of the certain compounds in the examples.
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a)

## Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

International application No.  
PCT/IB2004/000401

Patent document cited in search report	Publication data	Patent family member	Publication data
WO0116106 A1	08-03-2001	JP2001081031 A	27-03-2001
		AU200068416 A	26-03-2001
		BR200013648 A	07-05-2002
		NO200200952 A	20-03-2002
		EP1208086 A1	29-05-2002
		CZ200200724 A3	17-07-2002
		KR2002023424 A	28-03-2002
		SK200200261 A3	10-09-2002
		CN1371366 A	25-09-2002
		ZA200201424 A	27-11-2002
		JP2003508386T T	04-03-2003
		HU200203330 A2	28-02-2003
		NZ517520 A	30-05-2003
		US6638530 B1	28-10-2003
		MX2002002090 A1	01-08-2003
EP0847792 A1	17-06-1998	DE69729364E	08-07-2004
		US5769928 A	23-06-1998
		CA2224471 A	12-06-1998
		CN1195571 A	14-10-1998
		KR98063917 A	07-10-1998
		BR9706248 A	10-08-1999
		CA2224471 C	10-10-2000
		KR322957 B	08-03-2002
		EP0847792 B1	02-06-2004